

COUPLED PAINLEVÉ III SYSTEM WITH AFFINE WEYL GROUP SYMMETRY OF TYPE $D_6^{(1)}$

YUSUKE SASANO

ABSTRACT. We find and study a six-parameter family of coupled Painlevé III systems in dimension six with affine Weyl group symmetry of type $D_6^{(1)}$. We also find and study its degenerate systems with affine Weyl group symmetry of types $B_5^{(1)}$ and $D_5^{(2)}$.

1. INTRODUCTION

In [13, 14], we presented a 4-parameter family of 2-coupled Painlevé III systems in dimension four with affine Weyl group symmetry of type $D_4^{(1)}$. We will make non-linear ordinary differential systems with affine Weyl group symmetry of type $D_{2n+2}^{(1)}$ ($n \geq 2$).

In [10, 12], we succeeded to make $(2n+2)$ -parameter family of n -coupled Painlevé VI systems in dimension $2n$ with affine Weyl group symmetry of type $D_{2n+2}^{(1)}$ ($n \geq 1$) by connecting the invariant divisors $p_i, q_i - q_{i+1}, p_{i+1}$ for the canonical variables (q_i, p_i) ($i = 1, 2, \dots, n$). These systems are polynomial Hamiltonian systems with coupled Painlevé VI Hamiltonians given by

$$(1) \quad \frac{dq_i}{dt} = \frac{\partial H}{\partial p_i}, \quad \frac{dp_i}{dt} = -\frac{\partial H}{\partial q_i} \quad (i = 1, 2, \dots, n)$$

with the polynomial Hamiltonian

$$(2) \quad H = \sum_{i=1}^n H_{VI}(q_i, p_i, t; \alpha_0^{(i)}, \alpha_1^{(i)}, \dots, \alpha_4^{(i)}) + \sum_{1 \leq l < m \leq n} \frac{2(q_l - t)p_l q_m((q_m - 1)p_m + \alpha_2^{(m)})}{t(t-1)},$$

where the symbol $H_{VI}(x, y, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4)$ is given by

$$(3) \quad \begin{aligned} & H_{VI}(x, y, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4) \\ &= \frac{1}{t(t-1)} [y^2(x-t)(x-1)x - \{(\alpha_0 - 1)(x-1)x + \alpha_3(x-t)x \\ &+ \alpha_4(x-t)(x-1)\}y + \alpha_2(\alpha_1 + \alpha_2)x] \quad (\alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3 + \alpha_4 = 1). \end{aligned}$$

However, in this case the invariant divisors are different from the ones of P_{VI} -case.

Invariant divisors	f_0	f_1	f_2	f_3	f_4
P_{VI}	$q - t$	$q - \infty$	p	$q - 1$	q
2-CPIII	$p_1 - 1$	p_1	$q_1 q_2 - 1$	p_2	$p_2 - t$

2000 *Mathematics Subject Classification.* 34M55; 34M45; 58F05; 32S65.

Key words and phrases. Affine Weyl group, birational symmetry, coupled Painlevé system.

At first, let us consider $D_6^{(1)}$ case. In this paper, we present a 6-parameter family of coupled Painlevé III systems with affine Weyl group symmetry of type $D_6^{(1)}$ explicitly given by

$$(4) \quad \frac{dx}{dt} = \frac{\partial H}{\partial y}, \quad \frac{dy}{dt} = -\frac{\partial H}{\partial x}, \quad \frac{dz}{dt} = \frac{\partial H}{\partial w}, \quad \frac{dw}{dt} = -\frac{\partial H}{\partial z}, \quad \frac{dq}{dt} = \frac{\partial H}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial H}{\partial q}$$

with the polynomial Hamiltonian

$$(5) \quad \begin{aligned} H &= \frac{x^2(y-1)y + x\{(\alpha_0 + \alpha_1)y - \alpha_1\} + ty}{t} \\ &+ \frac{z^2(w-1)w + z\{(\alpha_0 + \alpha_1 + 2\alpha_2 + 2\alpha_3)w - \alpha_3\} + tw}{t} \\ &+ \frac{q^2(p-t)p + q\{(\alpha_5 + \alpha_6 - 1)p - t\alpha_5\} + p}{t} + \frac{2yz(zw + \alpha_3)}{t} - \frac{2(y+w)p}{t} \\ &= H_{III}(x, y, t; \alpha_1, \alpha_0) + H_{III}(z, w, t; \alpha_3, \alpha_0 + \alpha_1 + 2\alpha_2 + \alpha_3) \\ &+ \tilde{H}_{III}(q, p, t; \alpha_5, 1 - \alpha_6) + \frac{2yz(zw + \alpha_3)}{t} - \frac{2(y+w)p}{t}. \end{aligned}$$

Here x, y, z, w, q and p denote unknown complex variables, and $\alpha_0, \alpha_1, \dots, \alpha_6$ are complex parameters satisfying the relation:

$$(6) \quad \alpha_0 + \alpha_1 + 2\alpha_2 + 2\alpha_3 + 2\alpha_4 + \alpha_5 + \alpha_6 = 1.$$

The symbols H_{III}, \tilde{H}_{III} are given by

$$(7) \quad H_{III}(u, v, t; \gamma_0, \gamma_1, \gamma_2) = \frac{u^2v(v-1) + u\{(\gamma_0 + \gamma_2)v - \gamma_0\} + tv}{t} \quad (\gamma_0 + 2\gamma_1 + \gamma_2 = 1),$$

$$(8) \quad \tilde{H}_{III}(U, V, t; \gamma_0, \gamma_1, \gamma_2) = \frac{U^2V(V-t) - U\{(-\gamma_0 + \gamma_2)V + \gamma_0t\} + V}{t}.$$

The relation between (u, v) and (U, V) is given by

$$(9) \quad (U, V) = (1/u, -u(vu + \gamma_0)).$$

We remark that for this system we tried to seek its first integrals of polynomial type with respect to x, y, z, w, q, p . However, we can not find. Of course, the Hamiltonian H is not the first integral.

This is the second example which gave higher order Painlevé type systems of type $D_6^{(1)}$.

We also find and study its degenerate systems with affine Weyl group symmetry of types $B_5^{(1)}$ and $D_5^{(2)}$. In $D_5^{(2)}$ -case, each differential system with respect to all principal parts has its first integral. Nevertheless, the polynomial Hamiltonian itself is not its first integral (see Section 4).

We give an explicit confluence process from the $D_6^{(1)}$ system, respectively. We will show that the system of type $B_5^{(1)}$ is equivalent to the $D_5^{(1)}$ system (see [15]) by explicit birational and symplectic transformations with some parameter's changes (see Section 3).

2. THE SYSTEM OF TYPE $D_6^{(1)}$

In this section, we present a 6-parameter family of coupled Painlevé III systems with affine Weyl group symmetry of type $D_6^{(1)}$ explicitly given by

$$(10) \quad \left\{ \begin{array}{l} \frac{dx}{dt} = \frac{\partial H}{\partial y} = \frac{2x^2y + 2z^2w - x^2 - (\alpha_0 + \alpha_1)x + 2\alpha_3z - 2p + t}{t}, \\ \frac{dy}{dt} = -\frac{\partial H}{\partial x} = \frac{-2xy^2 + 2xy - (\alpha_0 + \alpha_1)y + \alpha_1}{t}, \\ \frac{dz}{dt} = \frac{\partial H}{\partial w} = \frac{2z^2w + 2yz^2 - z^2 - (2\alpha_4 - 1 + \alpha_5 + \alpha_6)z - 2p + t}{t}, \\ \frac{dw}{dt} = -\frac{\partial H}{\partial z} = \frac{-2zw^2 - 4yzw + 2zw - 2\alpha_3y + (2\alpha_4 - 1 + \alpha_5 + \alpha_6)w + \alpha_3}{t}, \\ \frac{dq}{dt} = \frac{\partial H}{\partial p} = \frac{2q^2p - tq^2 - 2y - 2w + (\alpha_5 + \alpha_6 - 1)q + 1}{t}, \\ \frac{dp}{dt} = -\frac{\partial H}{\partial q} = \frac{-2qp^2 + 2tqp - (\alpha_5 + \alpha_6 - 1)p + t\alpha_5}{t} \end{array} \right.$$

with the Hamiltonian (5).

THEOREM 2.1. *The system (10) admits extended affine Weyl group symmetry of type $D_6^{(1)}$ as the group of its Bäcklund transformations whose generators are explicitly given as follows: with the notation $(*) := (x, y, z, w, q, p, t; \alpha_0, \alpha_1, \dots, \alpha_6)$,*

$$\begin{aligned} s_0 : (*) &\rightarrow \left(x + \frac{\alpha_0}{y-1}, y, z, w, q, p, t; -\alpha_0, \alpha_1, \alpha_2 + \alpha_0, \alpha_3, \alpha_4, \alpha_5, \alpha_6 \right), \\ s_1 : (*) &\rightarrow \left(x + \frac{\alpha_1}{y}, y, z, w, q, p, t; \alpha_0, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4, \alpha_5, \alpha_6 \right), \\ s_2 : (*) &\rightarrow \left(x, y - \frac{\alpha_2}{x-z}, z, w + \frac{\alpha_2}{x-z}, q, p, t; \alpha_0 + \alpha_2, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4, \alpha_5, \alpha_6 \right), \\ s_3 : (*) &\rightarrow \left(x, y, z + \frac{\alpha_3}{w}, w, q, p, t; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4 + \alpha_3, \alpha_5, \alpha_6 \right), \\ s_4 : (*) &\rightarrow \left(x, y, z, w - \frac{\alpha_4q}{zq-1}, q, p - \frac{\alpha_4z}{zq-1}, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3 + \alpha_4, -\alpha_4, \alpha_5 + \alpha_4, \alpha_6 + \alpha_4 \right), \\ s_5 : (*) &\rightarrow \left(x, y, z, w, q + \frac{\alpha_5}{p}, p, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4 + \alpha_5, -\alpha_5, \alpha_6 \right), \\ s_6 : (*) &\rightarrow \left(x, y, z, w, q + \frac{\alpha_6}{p-t}, p, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4 + \alpha_6, \alpha_5, -\alpha_6 \right), \\ \pi_1 : (*) &\rightarrow (-x, 1-y, -z, -w, -q, t-p, t; \alpha_1, \alpha_0, \alpha_2, \alpha_3, \alpha_4, \alpha_6, \alpha_5), \\ \pi_2 : (*) &\rightarrow \left(tq, \frac{p}{t}, \frac{t}{z}, -\frac{(zw + \alpha_3)z}{t}, \frac{x}{t}, ty, t; \alpha_6, \alpha_5, \alpha_4, \alpha_3, \alpha_2, \alpha_1, \alpha_0 \right), \end{aligned}$$




FIGURE 1. The figure denotes the Dynkin diagram of types $B_4^{(1)}$, $D_4^{(1)}$ and $D_6^{(1)}$. The symbol in each circle denotes the invariant divisors of the systems of types $B_4^{(1)}$, $D_4^{(1)}$ and $D_6^{(1)}$.

$$\pi_3 : (*) \rightarrow (x, y, z, w, q, p-t, -t; \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_4, \alpha_6, \alpha_5),$$

$$\pi_4 : (*) \rightarrow (-x, 1-y, -z, -w, -q, -p, -t; \alpha_1, \alpha_0, \alpha_2, \alpha_3, \alpha_4, \alpha_5, \alpha_6).$$

The Bäcklund transformations of this system satisfy the universal description for $D_6^{(1)}$ root system. Since these universal Bäcklund transformations have Lie theoretic origin, similarity reduction of a Drinfeld-Sokolov hierarchy admits such a Bäcklund symmetry.

PROPOSITION 2.2. *Let us define the following translation operators:*

$$(11) \quad \begin{aligned} T_1 &:= \pi_1 s_5 s_4 s_3 s_2 s_1 s_0 s_1 s_2 s_3 s_4 s_5, & T_2 &:= s_4 s_6 T_1 s_6 s_4, \\ T_3 &:= s_6 T_1 s_6, & T_4 &:= \pi_2 T_1 \pi_2, & T_5 &:= \pi_2 T_2 \pi_2, & T_6 &:= \pi_2 T_3 \pi_2. \end{aligned}$$

These translation operators act on parameters α_i as follows:

$$(12) \quad \begin{aligned} T_1(\alpha_0, \alpha_1, \dots, \alpha_6) &= (\alpha_0, \alpha_1, \dots, \alpha_6) + (0, 0, 0, 0, 0, -1, 1), \\ T_2(\alpha_0, \alpha_1, \dots, \alpha_6) &= (\alpha_0, \alpha_1, \dots, \alpha_6) + (0, 0, 0, 1, -1, 0, 0), \\ T_3(\alpha_0, \alpha_1, \dots, \alpha_6) &= (\alpha_0, \alpha_1, \dots, \alpha_6) + (0, 0, 0, 0, 1, -1, -1), \\ T_4(\alpha_0, \alpha_1, \dots, \alpha_6) &= (\alpha_0, \alpha_1, \dots, \alpha_6) + (1, -1, 0, 0, 0, 0, 0), \\ T_5(\alpha_0, \alpha_1, \dots, \alpha_6) &= (\alpha_0, \alpha_1, \dots, \alpha_6) + (0, 0, -1, 1, 0, 0, 0), \\ T_6(\alpha_0, \alpha_1, \dots, \alpha_6) &= (\alpha_0, \alpha_1, \dots, \alpha_6) + (-1, -1, 1, 0, 0, 0, 0). \end{aligned}$$

THEOREM 2.3. *Let us consider a polynomial Hamiltonian system with Hamiltonian $H \in \mathbb{C}(t)[x, y, z, w, q, p]$. We assume that*

(A1) $\deg(H) = 4$ with respect to x, y, z, w, q, p .

(A2) This system becomes again a polynomial Hamiltonian system in each coordinate system r_i ($i = 0, 1, 2, 3, 5, 6$):

$$\begin{aligned} r_0 : x_0 &= 1/x, \quad y_0 = -((y-1)x + \alpha_0)x, \quad z_0 = z, \quad w_0 = w, \quad q_0 = q, \quad p_0 = p, \\ r_1 : x_1 &= 1/x, \quad y_1 = -(yx + \alpha_1)x, \quad z_1 = z, \quad w_1 = w, \quad q_1 = q, \quad p_1 = p, \\ r_2 : x_2 &= -((x-z)y - \alpha_2)y, \quad y_2 = 1/y, \quad z_2 = z, \quad w_2 = w + y, \quad q_2 = q, \quad p_2 = p, \\ r_3 : x_3 &= x, \quad y_3 = y, \quad z_3 = 1/z, \quad w_3 = -(wz + \alpha_3)z, \quad q_3 = q, \quad p_3 = p, \\ r_5 : x_5 &= x, \quad y_5 = y, \quad z_5 = z, \quad w_5 = w, \quad q_5 = 1/q, \quad p_5 = -(pq + \alpha_5)q, \\ r_6 : x_6 &= x, \quad y_6 = y, \quad z_6 = z, \quad w_6 = w, \quad q_6 = 1/q, \quad p_6 = -((p-t)q + \alpha_6)q. \end{aligned}$$

(A3) In addition to the assumption (A2), the Hamiltonian system in the coordinate system r_3 becomes again a polynomial Hamiltonian system in the coordinate system r_4 :

$$r_4 : x_4 = x_3, \quad y_4 = y_3, \quad z_4 = -((z_3 - q_3)w_3 - \alpha_4)w_3, \quad w_4 = 1/w_3, \quad q_4 = q_3, \quad p_4 = p_3 + w_3.$$

Then such a system coincides with the system (10) with the polynomial Hamiltonian (5).

We note that the conditions (A2) and (A3) should be read that

$$r_j(H) \quad (j = 0, 1, 2, 3, 5), \quad r_6(H + q), \quad r_4(r_3(H))$$

are polynomials with respect to x, y, z, w, q, p or $x_3, y_3, z_3, w_3, q_3, p_3$.

Finally, we consider the rational and algebraic solutions of the system (10).

At first, we consider the Dynkin diagram automorphism π_1 . By this transformation, the fixed solution is derived from

$$(13) \quad \begin{aligned} \alpha_0 &= \alpha_1, \quad \alpha_1 = \alpha_0, \quad \alpha_5 = \alpha_6, \quad \alpha_6 = \alpha_5, \\ x &= -x, \quad y = 1 - y, \quad z = -z, \quad w = -w, \quad q = -q, \quad p = t - p. \end{aligned}$$

Then we obtain

$$(14) \quad \begin{aligned} (\alpha_0, \alpha_1, \dots, \alpha_6) &= \left(\frac{1}{2} - \alpha_2 - \alpha_3 - \alpha_4 - \alpha_6, \frac{1}{2} - \alpha_2 - \alpha_3 - \alpha_4 - \alpha_6, \alpha_2, \alpha_3, \alpha_4, \alpha_6, \alpha_6 \right), \\ (x, y, z, w, q, p) &= \left(0, \frac{1}{2}, 0, 0, 0, \frac{t}{2} \right). \end{aligned}$$

Next, we find two algebraic solutions:

$$(15) \quad \begin{aligned} (\alpha_0, \alpha_1, \dots, \alpha_6) &= \left(\frac{1 - 2\alpha_3}{2}, 0, 0, \alpha_3, 0, 0, \frac{1 - 2\alpha_3}{2} \right), \\ (x, y, z, w, q, p) &= \left(\sqrt{t}, 0, \sqrt{t}, -\frac{\alpha_3}{2\sqrt{t}}, \frac{1}{\sqrt{t}}, 0 \right) \end{aligned}$$

and

$$(16) \quad \begin{aligned} (\alpha_0, \alpha_1, \dots, \alpha_6) &= \left(0, \frac{1+2\alpha_3}{2}, -\alpha_3, \alpha_3, -\alpha_3, \frac{1+2\alpha_3}{2}, 0\right), \\ (x, y, z, w, q, p) &= \left(-\sqrt{t}, 1, \sqrt{t}, -\frac{\alpha_3}{2\sqrt{t}}, -\frac{1}{\sqrt{t}}, t\right). \end{aligned}$$

3. THE SYSTEM OF TYPE $B_5^{(1)}$

In this section, we present a 5-parameter family of coupled Painlevé systems with affine Weyl group symmetry of type $B_5^{(1)}$ explicitly given by

$$(17) \quad \left\{ \begin{aligned} \frac{dx}{dt} &= \frac{\partial H}{\partial y} = \frac{2x^2y + 2z^2w + 2\alpha_0x + 2\alpha_2z - 2p + t}{t}, \\ \frac{dy}{dt} &= -\frac{\partial H}{\partial x} = \frac{-2xy^2 - 2\alpha_0y - 1}{t}, \\ \frac{dz}{dt} &= \frac{\partial H}{\partial w} = \frac{2z^2w + 2yz^2 + 2(\alpha_0 + \alpha_1 + \alpha_2)z - 2p + t}{t}, \\ \frac{dw}{dt} &= -\frac{\partial H}{\partial z} = \frac{-2zw^2 - 4yzw - 2\alpha_2y - 2(\alpha_0 + \alpha_1 + \alpha_2)w}{t}, \\ \frac{dq}{dt} &= \frac{\partial H}{\partial p} = \frac{2q^2p - tq^2 - 2y - 2w - 2(\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3)q}{t}, \\ \frac{dp}{dt} &= -\frac{\partial H}{\partial q} = \frac{-2qp^2 + 2tqp + 2(\alpha_0 + \alpha_1 + \alpha_2 + \alpha_3)p + t\alpha_4}{t} \end{aligned} \right.$$

with the polynomial Hamiltonian

$$(18) \quad \begin{aligned} H &= \frac{x^2y^2 + 2\alpha_0xy + x + ty}{t} + \frac{z^2w^2 + 2(\alpha_0 + \alpha_1 + \alpha_2)zw + tw}{t} \\ &+ \frac{q^2p^2 - tq^2p + (\alpha_4 + \alpha_5 - 1)qp - \alpha_4tq}{t} + \frac{2yz(zw + \alpha_2)}{t} - \frac{2(y+w)p}{t} \\ &= H_{III}^{D_7^{(1)}}(x, y, t; 2\alpha_0) + H_1(z, w, t; 2(\alpha_0 + \alpha_1 + \alpha_2)) \\ &+ H_2(q, p, t; \alpha_4 + \alpha_5 - 1, -\alpha_4) + \frac{2yz(zw + \alpha_2)}{t} - \frac{2(y+w)p}{t}. \end{aligned}$$

Here x, y, z, w, q and p denote unknown complex variables, and $\alpha_0, \alpha_1, \dots, \alpha_5$ are complex parameters satisfying the relation:

$$(19) \quad 2\alpha_0 + 2\alpha_1 + 2\alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5 = 1.$$

The symbols $H_{III}^{D_7^{(1)}}$, H_1 and H_2 are given by

$$(20) \quad H_{III}^{D_7^{(1)}}(q, p, t; \beta_1) = \frac{q^2 p^2 + \beta_1 q p + q + t p}{t} \quad (\beta_0 + \beta_1 = 1),$$

$$(21) \quad H_1(q, p, t; \alpha) = \frac{q^2 p^2 + \alpha q p + t p}{t},$$

$$(22) \quad H_2(q, p, t; \alpha, \beta) = \frac{q^2 p^2 - t q^2 p + \alpha q p + \beta t q}{t}.$$

We remark that for $y = q/\tau$, $t = \tau^2$ the Hamiltonian system with $H_{III}^{D_7^{(1)}}$ is the special case of the third Painlevé system (see [17]):

$$(23) \quad \frac{d^2 y}{d\tau^2} = \frac{1}{y} \left(\frac{dy}{d\tau} \right)^2 - \frac{1}{\tau} \frac{dy}{d\tau} + \frac{1}{\tau} (a y^2 + b) + c y^3 + \frac{d}{y}$$

with

$$(24) \quad a = -8, \quad b = 4(1 - \beta_1), \quad c = 0, \quad d = -4.$$

From the viewpoint of symmetry, the Hamiltonian system

$$(25) \quad \frac{dq}{dt} = \frac{\partial H_{III}^{D_7^{(1)}}}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial H_{III}^{D_7^{(1)}}}{\partial q}$$

has extended affine Weyl group symmetry of type $A_1^{(1)}$, whose generators $< s_0, s_1, \pi = \sigma \circ s_1 >$ are explicitly given as follows (see [17]):

$$(26) \quad \begin{cases} s_0(q, p, t; \beta_0, \beta_1) = \left(q, p + \frac{\beta_0}{q} - \frac{t}{q^2}, -t; -\beta_0, \beta_1 + 2\beta_0 \right), \\ s_1(q, p, t; \beta_0, \beta_1) = \left(-q + \frac{\beta_1}{p} + \frac{1}{p^2}, -p, -t; \beta_0 + 2\beta_1, -\beta_1 \right), \\ \sigma(q, p, t; \beta_0, \beta_1) = \left(t p, -\frac{q}{t}, -t; \beta_1, \beta_0 \right). \end{cases}$$

PROPOSITION 3.1. *By the following birational and symplectic transformations tr_i ($i = 1, 2$):*

$$(27) \quad \begin{cases} tr_1(q, p) = (q/t, t p), \\ tr_2(q, p) = (-p/t, t q), \end{cases}$$

the Hamiltonians H_1 and H_2 satisfy the following relations:

$$(28) \quad tr_1(H_1) = \frac{q^2 p^2 + (\alpha - 1) q p + p}{t}, \quad tr_2(H_2) = \frac{q^2 p^2 + q p^2 - (\alpha + 1) q p + \beta p}{t}.$$

Here, for notational convenience, we use the same symbol q, p, α .

By Proposition 3.1, we see that the Hamiltonian system with

$$K_1 := \frac{q^2 p^2 + (\alpha - 1) q p + p}{t}$$

has the first integral.

PROPOSITION 3.2. *The system with the Hamiltonian K_1*

$$(29) \quad \frac{dq}{dt} = \frac{\partial K_1}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial K_1}{\partial q}$$

has the first integral I_1 :

$$(30) \quad I_1 = q^2 p^2 + (\alpha - 1)qp + p.$$

We see that the relation between the Hamiltonian K_1 and the first integral I_1 is explicitly given by

$$(31) \quad tK_1 = I_1.$$

We also show that the Hamiltonian system with

$$K_2 := \frac{q^2 p^2 + qp^2 - (\alpha + 1)qp + \beta p}{t}$$

has the first integral.

PROPOSITION 3.3. *The system with the Hamiltonian K_2*

$$(32) \quad \frac{dq}{dt} = \frac{\partial K_2}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial K_2}{\partial q}$$

has the first integral I_2 :

$$(33) \quad I_2 = q^2 p^2 + qp^2 - (\alpha + 1)qp + \beta p.$$

We see that the relation between the Hamiltonian K_2 and the first integral I_2 is explicitly given by

$$(34) \quad tK_2 = I_2.$$

We remark that for this system we tried to seek its first integrals of polynomial type with respect to x, y, z, w, q, p . However, we can not find. Of course, the Hamiltonian H is not the first integral.

THEOREM 3.4. *The system (17) admits extended affine Weyl group symmetry of type $B_5^{(1)}$ as the group of its Bäcklund transformations, whose generators are explicitly given*

as follows: with the notation $(*) := (x, y, z, w, q, p, t; \alpha_0, \alpha_1, \dots, \alpha_5)$,

$$\begin{aligned}
s_0 : (*) &\rightarrow \left(-x - \frac{2\alpha_0}{y} - \frac{1}{y^2}, -y, -z, -w, -q, -p, -t; -\alpha_0, \alpha_1 + 2\alpha_0, \alpha_2, \alpha_3, \alpha_4, \alpha_5 \right), \\
s_1 : (*) &\rightarrow \left(x, y - \frac{\alpha_1}{x-z}, z, w + \frac{\alpha_1}{x-z}, q, p, t; \alpha_0 + \alpha_1, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4, \alpha_5 \right), \\
s_2 : (*) &\rightarrow \left(x, y, z + \frac{\alpha_2}{w}, w, q, p, t; \alpha_0, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4, \alpha_5 \right), \\
s_3 : (*) &\rightarrow \left(x, y, z, w - \frac{\alpha_3 q}{zq-1}, q, p - \frac{\alpha_3 z}{zq-1}, t; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4 + \alpha_3, \alpha_5 + \alpha_3 \right), \\
s_4 : (*) &\rightarrow \left(x, y, z, w, q + \frac{\alpha_4}{p}, p, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3 + \alpha_4, -\alpha_4, \alpha_5 \right), \\
s_5 : (*) &\rightarrow \left(x, y, z, w, q + \frac{\alpha_5}{p-t}, p, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3 + \alpha_5, \alpha_4, -\alpha_5 \right), \\
\pi : (*) &\rightarrow (x, y, z, w, q, p-t, -t; \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_4).
\end{aligned}$$

PROPOSITION 3.5. *Let us define the following translation operators:*

$$\begin{aligned}
(35) \quad T_1 &:= \pi s_4 s_3 s_2 s_1 s_0 s_1 s_2 s_3 s_4, \quad T_2 := \pi s_5 s_4 s_3 s_2 s_1 s_0 s_1 s_2 s_3, \\
T_3 &:= s_3 s_5 T_1 s_5 s_3, \quad T_4 := s_2 T_3 s_2, \quad T_5 := s_1 T_4 s_1.
\end{aligned}$$

These translation operators act on parameters α_i as follows:

$$\begin{aligned}
(36) \quad T_1(\alpha_0, \alpha_1, \dots, \alpha_5) &= (\alpha_0, \alpha_1, \dots, \alpha_5) + (0, 0, 0, 0, -1, 1), \\
T_2(\alpha_0, \alpha_1, \dots, \alpha_5) &= (\alpha_0, \alpha_1, \dots, \alpha_5) + (0, 0, 0, -1, 1, 1), \\
T_3(\alpha_0, \alpha_1, \dots, \alpha_5) &= (\alpha_0, \alpha_1, \dots, \alpha_5) + (0, 0, 1, -1, 0, 0), \\
T_4(\alpha_0, \alpha_1, \dots, \alpha_5) &= (\alpha_0, \alpha_1, \dots, \alpha_5) + (0, 1, -1, 0, 0, 0), \\
T_5(\alpha_0, \alpha_1, \dots, \alpha_5) &= (\alpha_0, \alpha_1, \dots, \alpha_5) + (1, -1, 0, 0, 0, 0).
\end{aligned}$$

THEOREM 3.6. *Let us consider a polynomial Hamiltonian system with Hamiltonian $H \in \mathbb{C}(t)[x, y, z, w, q, p]$. We assume that*

(B1) *$\deg(H) = 4$ with respect to x, y, z, w, q, p .*

(B2) *This system becomes again a polynomial Hamiltonian system in each coordinate system r_i ($i = 0, 1, 2, 4, 5$):*

$$\begin{aligned}
r_0 : x_0 &= x + \frac{2\alpha_0}{y} + \frac{1}{y^2}, \quad y_0 = y, \quad z_0 = z, \quad w_0 = w, \quad q_0 = q, \quad p_0 = p, \\
r_1 : x_1 &= -((x-z)y - \alpha_1)y, \quad y_1 = 1/y, \quad z_1 = z, \quad w_1 = w + y, \quad q_1 = q, \quad p_1 = p, \\
r_2 : x_2 &= x, \quad y_2 = y, \quad z_2 = 1/z, \quad w_2 = -(wz + \alpha_2)z, \quad q_2 = q, \quad p_2 = p, \\
r_4 : x_4 &= x, \quad y_4 = y, \quad z_4 = z, \quad w_4 = w, \quad q_4 = 1/q, \quad p_4 = -(pq + \alpha_4)q, \\
r_5 : x_5 &= x, \quad y_5 = y, \quad z_5 = z, \quad w_5 = w, \quad q_5 = 1/q, \quad p_5 = -((p-t)q + \alpha_5)q.
\end{aligned}$$

(B3) In addition to the assumption (B2), the Hamiltonian system in the coordinate system $(x_4, y_4, z_4, w_4, q_4, p_4)$ becomes again a polynomial Hamiltonian system in the coordinate system r_3 :

$$r_3 : x_3 = x_4, \quad y_3 = y_4, \quad z_3 = -((z_4 - q_4)w_4 - \alpha_3)w_4, \quad w_3 = 1/w_4, \quad q_3 = q_4, \quad p_3 = p_4 + w_4.$$

Then such a system coincides with the system (17) with the polynomial Hamiltonian (18).

We note that the conditions (B2) and (B3) should be read that

$$r_j(H) \quad (j = 0, 1, 2, 4), \quad r_5(H + q), \quad r_3(r_4(H))$$

are polynomials with respect to x, y, z, w, q, p or $x_4, y_4, z_4, w_4, q_4, p_4$.

Theorems 3.4 and 3.6 can be checked by a direct calculation, respectively.

Next, we show the confluence process from the system (10) to the system (17).

THEOREM 3.7. *For the system (10) of type $D_6^{(1)}$, we make the change of parameters and variables*

(37)

$$\alpha_0 = \frac{1}{\varepsilon} + 2A_0, \quad \alpha_1 = -\frac{1}{\varepsilon}, \quad \alpha_2 = A_1, \quad \alpha_3 = A_2, \quad \alpha_4 = A_3, \quad \alpha_5 = A_4, \quad \alpha_6 = A_5,$$

$$(38) \quad t = \varepsilon T, \quad x = \varepsilon X, \quad y = \frac{Y}{\varepsilon}, \quad z = \varepsilon Z, \quad w = \frac{W}{\varepsilon}, \quad q = \frac{Q}{\varepsilon}, \quad p = \varepsilon P$$

from $\alpha_0, \alpha_1, \dots, \alpha_6, x, y, z, w, q, p$ to $A_0, A_1, \dots, A_5, X, Y, Z, W, Q, P$. Then the system (10) can also be written in the new variables X, Y, Z, W, Q, P and parameters A_0, A_1, \dots, A_5 as a Hamiltonian system. This new system tends to the system (17) with the Hamiltonian (18) as $\varepsilon \rightarrow 0$.

Finally, we show the relation between the system (17) and the system of type $D_5^{(1)}$ (see [15]).

THEOREM 3.8. *For the system (17) of type $B_5^{(1)}$, we make the change of parameters and variables*

$$(39) \quad \alpha_0 = \frac{A_0 - A_1}{2}, \quad \alpha_1 = A_1, \quad \alpha_2 = A_2, \quad \alpha_3 = A_3, \quad \alpha_4 = A_4, \quad \alpha_5 = A_5,$$

$$(40) \quad X = ((x - z)y - \alpha_1)y, \quad Y = -1/y, \quad Z = z, \quad W = w + y, \quad Q = q, \quad P = p$$

from $\alpha_0, \alpha_1, \dots, \alpha_5, x, y, z, w, q, p$ to $A_0, A_1, \dots, A_5, X, Y, Z, W, Q, P$. Then the system (10) can also be written in the new variables X, Y, Z, W, Q, P and parameters A_0, A_1, \dots, A_5 as a Hamiltonian system. This new system tends to

$$(41) \quad \frac{dx}{dt} = \frac{\partial H}{\partial y}, \quad \frac{dy}{dt} = -\frac{\partial H}{\partial x}, \quad \frac{dz}{dt} = \frac{\partial H}{\partial w}, \quad \frac{dw}{dt} = -\frac{\partial H}{\partial z}, \quad \frac{dq}{dt} = \frac{\partial H}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial H}{\partial q}$$

with the polynomial Hamiltonian

$$(42) \quad H = \frac{x^2y^2 + xy^2 - (\alpha_0 + \alpha_1)xy - \alpha_0y}{t} + \frac{z^2w^2 + (\alpha_0 + \alpha_1 + 2\alpha_2)zw + z + tw}{t} \\ + \frac{q^2p^2 - tq^2p - (1 - \alpha_4 - \alpha_5)qp - \alpha_4tq}{t} + \frac{2(xz - wp)}{t} \\ (\alpha_0 + \alpha_1 + 2\alpha_2 + 2\alpha_3 + \alpha_4 + \alpha_5 = 1).$$

Here, for notational convenience, we have renamed A_i, X, Y, Z, W, Q, P to $\alpha_i, x, y, z, w, q, p$ (which are not the same as the previous $\alpha_i, x, y, z, w, q, p$).

We recall that this system admits extended affine Weyl group symmetry of type $D_5^{(1)}$ (see [15]) as the group of its Bäcklund transformations whose generators are explicitly given as follows: with the notation $(*) := (x, y, z, w, q, p, t; \alpha_0, \alpha_1, \dots, \alpha_5)$,




FIGURE 2. Dynkin diagram of type $D_5^{(1)}$

$$s_0 : (*) \rightarrow \left(x, y - \frac{\alpha_0}{x}, z, w, q, p, t; -\alpha_0, \alpha_1, \alpha_2 + \alpha_0, \alpha_3, \alpha_4, \alpha_5 \right), \\ s_1 : (*) \rightarrow \left(x, y - \frac{\alpha_1}{x+1}, z, w, q, p, t; \alpha_0, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4, \alpha_5 \right), \\ s_2 : (*) \rightarrow \left(x + \frac{\alpha_2 w}{yw+1}, y, z + \frac{\alpha_2 y}{yw+1}, w, q, p, t; \alpha_0 + \alpha_2, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4, \alpha_5 \right), \\ s_3 : (*) \rightarrow \left(x, y, z, w - \frac{\alpha_3 q}{zq-1}, q, p - \frac{\alpha_3 z}{zq-1}, t; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4 + \alpha_3, \alpha_5 + \alpha_3 \right), \\ s_4 : (*) \rightarrow \left(x, y, z, w, q + \frac{\alpha_4}{p}, p, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3 + \alpha_4, -\alpha_4, \alpha_5 \right), \\ s_5 : (*) \rightarrow \left(x, y, z, w, q + \frac{\alpha_5}{p-t}, p, t; \alpha_0, \alpha_1, \alpha_2, \alpha_3 + \alpha_5, \alpha_4, -\alpha_5 \right),$$

$$\begin{aligned}
\pi_1 : (*) &\rightarrow (-x-1, -y, -z, -w, -q, -p, -t; \alpha_1, \alpha_0, \alpha_2, \alpha_3, \alpha_4, \alpha_5), \\
\pi_2 : (*) &\rightarrow (x, y, z, w, q, p-t, -t; \alpha_0, \alpha_1, \alpha_2, \alpha_3, \alpha_5, \alpha_4), \\
\pi_3 : (*) &\rightarrow \left(\frac{(p-t)}{t}, -tq, -tw, \frac{z}{t}, \frac{y}{t}, -t(x+1), -t; \alpha_5, \alpha_4, \alpha_3, \alpha_2, \alpha_1, \alpha_0 \right).
\end{aligned}$$

4. THE SYSTEM OF TYPE $D_5^{(2)}$

In this section, we present a 4-parameter family of coupled Hamiltonian systems in dimension six with extended affine Weyl group symmetry of type $D_5^{(2)}$ given by

$$(43) \quad \left\{ \begin{aligned} \frac{dx}{dt} &= \frac{\partial H}{\partial y} = \frac{x^2y + z^2w + \alpha_0x + \alpha_2z - p}{t}, \\ \frac{dy}{dt} &= -\frac{\partial H}{\partial x} = \frac{-2xy^2 - 2\alpha_0y - 1}{2t}, \\ \frac{dz}{dt} &= \frac{\partial H}{\partial w} = \frac{z^2w + yz^2 + (\alpha_0 + \alpha_1 + \alpha_2)z - p}{t}, \\ \frac{dw}{dt} &= -\frac{\partial H}{\partial z} = \frac{-zw^2 - 2yzw - \alpha_2y - (\alpha_0 + \alpha_1 + \alpha_2)w}{t}, \\ \frac{dq}{dt} &= \frac{\partial H}{\partial p} = \frac{q^2p - y - w + (\alpha_4 - 1)q}{t}, \\ \frac{dp}{dt} &= -\frac{\partial H}{\partial q} = \frac{-2qp^2 - 2(\alpha_4 - 1)p + t}{2t} \end{aligned} \right.$$

with the polynomial Hamiltonian

$$\begin{aligned}
(44) \quad H &= \frac{x^2y^2 + 2\alpha_0xy + x}{2t} + \frac{z^2w^2 + 2(\alpha_0 + \alpha_1 + \alpha_2)zw}{2t} \\
&+ \frac{q^2p^2 + 2(\alpha_4 - 1)qp - tq}{2t} + \frac{yz(zw + \alpha_2)}{t} - \frac{(y+w)p}{t} \\
&= H_3(x, y, t; 2\alpha_0) + H_4(z, w, t; 2(\alpha_0 + \alpha_1 + \alpha_2)) \\
&+ H_5(q, p, t; 2(\alpha_4 - 1)) + \frac{yz(zw + \alpha_2)}{t} - \frac{(y+w)p}{t}.
\end{aligned}$$

Here x, y, z, w, q and p denote unknown complex variables and $\alpha_0, \alpha_1, \dots, \alpha_4$ are complex parameters satisfying the relation:

$$(45) \quad \alpha_0 + \alpha_1 + \alpha_2 + \alpha_3 + \alpha_4 = 1.$$

The symbols H_3, H_4 and H_5 are given by

$$(46) \quad H_3(q, p, t; \alpha) = \frac{q^2p^2 + \alpha qp + q}{2t},$$

$$(47) \quad H_4(q, p, t; \alpha) = \frac{q^2p^2 + \alpha qp}{2t},$$

$$(48) \quad H_5(q, p, t; \alpha, \beta) = \frac{q^2p^2 + \alpha qp - tq}{2t}.$$

PROPOSITION 4.1. *The system with the Hamiltonian H_3*

$$(49) \quad \frac{dq}{dt} = \frac{\partial H_3}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial H_3}{\partial q}$$

has the first integral I_3 :

$$(50) \quad I_3 = q^2 p^2 + \alpha q p + q.$$

PROPOSITION 4.2. *The system with the Hamiltonian H_4*

$$(51) \quad \frac{dq}{dt} = \frac{\partial H_4}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial H_4}{\partial q}$$

has the first integral I_4 :

$$(52) \quad I_4 = qp.$$

PROPOSITION 4.3. *By the following birational and symplectic transformation tr_5 :*

$$(53) \quad tr_5(q, p) = (tq, p/t),$$

the Hamiltonians H_5 satisfy the following relation:

$$(54) \quad tr_5(H_5) = \frac{q^2 p^2 + \alpha q p - q}{2t}.$$

By Proposition 4.3, we see that the Hamiltonian system with

$$K_5 := \frac{q^2 p^2 + \alpha q p - q}{2t}$$

has the first integral.

PROPOSITION 4.4. *The system with the Hamiltonian K_5*

$$(55) \quad \frac{dq}{dt} = \frac{\partial K_5}{\partial p}, \quad \frac{dp}{dt} = -\frac{\partial K_5}{\partial q}$$

has the first integral I_5 :

$$(56) \quad I_5 = q^2 p^2 + \alpha q p - q.$$

In this case, each differential system with respect to all principal parts H_3, H_4 and H_5 has its first integral. Nevertheless, the Hamiltonian H is not the first integral. For this system we tried to seek its first integrals of polynomial type with respect to x, y, z, w, q, p . However, we can not find.

THEOREM 4.5. *The system (43) admits extended affine Weyl group symmetry of type $D_5^{(2)}$ as the group of its Bäcklund transformations, whose generators are explicitly given*

as follows: with the notation $(*) := (x, y, z, w, q, p, t; \alpha_0, \alpha_1, \dots, \alpha_4)$,

$$\begin{aligned}
s_0 : (*) &\rightarrow \left(-x - \frac{2\alpha_0}{y} - \frac{1}{y^2}, -y, -z, -w, -q, -p, -t; -\alpha_0, \alpha_1 + 2\alpha_0, \alpha_2, \alpha_3, \alpha_4 \right), \\
s_1 : (*) &\rightarrow \left(x, y - \frac{\alpha_1}{x-z}, z, w + \frac{\alpha_1}{x-z}, q, p, t; \alpha_0 + \alpha_1, -\alpha_1, \alpha_2 + \alpha_1, \alpha_3, \alpha_4 \right), \\
s_2 : (*) &\rightarrow \left(x, y, z + \frac{\alpha_2}{w}, w, q, p, t; \alpha_0, \alpha_1 + \alpha_2, -\alpha_2, \alpha_3 + \alpha_2, \alpha_4 \right), \\
s_3 : (*) &\rightarrow \left(x, y, z, w - \frac{\alpha_3 q}{zq-1}, q, p - \frac{\alpha_3 z}{zq-1}, t; \alpha_0, \alpha_1, \alpha_2 + \alpha_3, -\alpha_3, \alpha_4 + \alpha_3 \right), \\
s_4 : (*) &\rightarrow \left(x, y, z, w, q + \frac{2\alpha_4}{p} - \frac{t}{p^2}, p, -t; \alpha_0, \alpha_1, \alpha_2, \alpha_3 + 2\alpha_4, -\alpha_4 \right), \\
\pi : (*) &\rightarrow \left(-tq, -\frac{p}{t}, -\frac{t}{z}, \frac{(zw + \alpha_2)z}{t}, -\frac{x}{t}, -ty, t; \alpha_4, \alpha_3, \alpha_2, \alpha_1, \alpha_0 \right).
\end{aligned}$$

PROPOSITION 4.6. *Let us define the following translation operators:*

$$\begin{aligned}
(57) \quad T_1 &:= s_4 s_3 s_2 s_1 s_0 s_1 s_2 s_3, \quad T_2 := s_3 T_1 s_3, \\
T_3 &:= s_2 T_2 s_2, \quad T_4 := s_1 T_3 s_1.
\end{aligned}$$

These translation operators act on parameters α_i as follows:

$$\begin{aligned}
(58) \quad T_1(\alpha_0, \alpha_1, \dots, \alpha_4) &= (\alpha_0, \alpha_1, \dots, \alpha_4) + (0, 0, 0, -2, 2), \\
T_2(\alpha_0, \alpha_1, \dots, \alpha_4) &= (\alpha_0, \alpha_1, \dots, \alpha_4) + (0, 0, -2, 2, 0), \\
T_3(\alpha_0, \alpha_1, \dots, \alpha_4) &= (\alpha_0, \alpha_1, \dots, \alpha_4) + (0, -2, 2, 0, 0), \\
T_4(\alpha_0, \alpha_1, \dots, \alpha_4) &= (\alpha_0, \alpha_1, \dots, \alpha_4) + (-2, 2, 0, 0, 0).
\end{aligned}$$

THEOREM 4.7. *Let us consider a polynomial Hamiltonian system with Hamiltonian $H \in \mathbb{C}(t)[x, y, z, w, q, p]$. We assume that*

(C1) *$\deg(H) = 4$ with respect to x, y, z, w, q, p .*

(C2) *This system becomes again a polynomial Hamiltonian system in each coordinate system r_i ($i = 0, 1, 2, 4$):*

$$\begin{aligned}
r_0 : x_0 &= x + \frac{2\alpha_0}{y} + \frac{1}{y^2}, \quad y_0 = y, \quad z_0 = z, \quad w_0 = w, \quad q_0 = q, \quad p_0 = p, \\
r_1 : x_1 &= -((x-z)y - \alpha_1)y, \quad y_1 = 1/y, \quad z_1 = z, \quad w_1 = w + y, \quad q_1 = q, \quad p_1 = p, \\
r_2 : x_2 &= x, \quad y_2 = y, \quad z_2 = 1/z, \quad w_2 = -(wz + \alpha_2)z, \quad q_2 = q, \quad p_2 = p, \\
r_4 : x_4 &= x, \quad y_4 = y, \quad z_4 = z, \quad w_4 = w, \quad q_4 = q + \frac{2\alpha_4}{p} - \frac{t}{p^2}, \quad p_4 = p.
\end{aligned}$$

(C3) *In addition to the assumption (C2), the Hamiltonian system in the coordinate system $(x_2, y_2, z_2, w_2, q_2, p_2)$ becomes again a polynomial Hamiltonian system in the coordinate system r_3 :*

$$r_3 : x_3 = x_2, \quad y_3 = y_2, \quad z_3 = -((z_2 - q_2)w_2 - \alpha_3)w_2, \quad w_3 = 1/w_2, \quad q_3 = q_2, \quad p_3 = p_2 + w_2.$$

Then such a system coincides with the system (43) with the polynomial Hamiltonian (44).

We note that the conditions (C2) and (C3) should be read that

$$r_j(H) \quad (j = 0, 1, 2), \quad r_4(H + 1/p), \quad r_3(r_2(H))$$

are polynomials with respect to x, y, z, w, q, p or $x_2, y_2, z_2, w_2, q_2, p_2$.

Theorems 4.5 and 4.7 can be checked by a direct calculation, respectively.

Next, we show the confluence process from the system (10) to the system (43).

THEOREM 4.8. *For the system (10) of type $D_6^{(1)}$, we make the change of parameters and variables*

(59)

$$\alpha_0 = -\frac{1}{\varepsilon} + A_0, \quad \alpha_1 = \frac{1}{\varepsilon}, \quad \alpha_2 = \frac{A_1}{2}, \quad \alpha_3 = \frac{A_2}{2}, \quad \alpha_4 = \frac{A_3}{2}, \quad \alpha_5 = \frac{1}{\varepsilon}, \quad \alpha_6 = -\frac{1}{\varepsilon} + A_4,$$

$$(60) \quad t = \frac{\varepsilon^2}{16}T, \quad x = \frac{\varepsilon}{4}X, \quad y = \frac{2Y}{\varepsilon}, \quad z = \frac{\varepsilon}{4}Z, \quad w = \frac{2W}{\varepsilon}, \quad q = \frac{4Q}{\varepsilon}, \quad p = \frac{\varepsilon}{8}P$$

from $\alpha_0, \alpha_1, \dots, \alpha_6, x, y, z, w, q, p$ to $A_0, A_1, \dots, A_4, X, Y, Z, W, Q, P$. Then the system (10) can also be written in the new variables X, Y, Z, W, Q, P and parameters A_0, A_1, \dots, A_4 as a Hamiltonian system. This new system tends to the system (43) with the Hamiltonian (44) as $\varepsilon \rightarrow 0$.

Finally, we find an algebraic solution of the system (43):

$$(61) \quad (\alpha_0, \alpha_1, \dots, \alpha_4) = (\alpha_0, \alpha_1, 1 - 2\alpha_0 - 2\alpha_1, \alpha_1, \alpha_0),$$

$$(x, y, z, w, q, p) = \left(-\left(\frac{1 + \sqrt{-1}}{4} \right) (t^{\frac{1}{4}} + 2(1 + \sqrt{-1})\sqrt{t}), -\frac{1 - \sqrt{-1}}{2t^{\frac{1}{4}}}, \sqrt{-t}, \right.$$

$$\left. -\frac{\sqrt{-1}(2\alpha_0 + 2\alpha_1 - 1)}{2\sqrt{t}}, \frac{(1 + \sqrt{-1}) + 4\sqrt{-1}t^{\frac{1}{4}}}{4t^{\frac{3}{4}}}, \left(\frac{1}{2} - \frac{\sqrt{-1}}{2} \right) t^{\frac{3}{4}} \right).$$

REFERENCES

- [1] P. Painlevé, *Mémoire sur les équations différentielles dont l'intégrale générale est uniforme*, Bull. Société Mathématique de France. **28** (1900), 201–261.
- [2] P. Painlevé, *Sur les équations différentielles du second ordre et d'ordre supérieur dont l'intégrale est uniforme*, Acta Math. **25** (1902), 1–85.
- [3] B. Gambier, *Sur les équations différentielles du second ordre et du premier degré dont l'intégrale générale est à points critiques fixes*, Acta Math. **33** (1910), 1–55.
- [4] C. M. Cosgrove and G. Scoufis, *Painlevé classification of a class of differential equations of the second order and second degree*, Studies in Applied Mathematics. **88** (1993), 25–87.
- [5] C. M. Cosgrove, *All binomial-type Painlevé equations of the second order and degree three or higher*, Studies in Applied Mathematics. **90** (1993), 119–187.
- [6] F. Bureau, *Integration of some nonlinear systems of ordinary differential equations*, Annali di Matematica. **94** (1972), 345–359.

- [7] J. Chazy, *Sur les équations différentielles dont l'intégrale générale est uniforme et admet des singularités essentielles mobiles*, Comptes Rendus de l'Académie des Sciences, Paris. **149** (1909), 563–565.
- [8] J. Chazy, *Sur les équations différentielles dont l'intégrale générale possède une coupure essentielle mobile*, Comptes Rendus de l'Académie des Sciences, Paris. **150** (1910), 456–458.
- [9] J. Chazy, *Sur les équations différentielles du troisième ordre et d'ordre supérieur dont l'intégrale a ses points critiques fixes*, Acta Math. **34** (1911), 317–385.
- [10] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of types $B_6^{(1)}$, $D_6^{(1)}$ and $D_7^{(2)}$* , preprint.
- [11] Y. Sasano, *Four-dimensional Painlevé systems of types $D_5^{(1)}$ and $B_4^{(1)}$* , preprint.
- [12] Y. Sasano, *Higher order Painlevé equations of type $D_l^{(1)}$* , RIMS Kokyuroku **1473** (2006), 143–163.
- [13] Y. Sasano, *Symmetries in the system of type $D_4^{(1)}$* , preprint.
- [14] Y. Sasano, *Coupled Painlevé III systems with affine Weyl group symmetry of types $B_4^{(1)}$, $D_4^{(1)}$ and $D_5^{(2)}$* , preprint.
- [15] Y. Sasano, *Coupled Painlevé III systems with affine Weyl group symmetry of types $B_5^{(1)}$, $D_5^{(1)}$ and $D_6^{(2)}$* , preprint.
- [16] Y. Sasano, *Coupled Painlevé VI systems in dimension four with affine Weyl group symmetry of type $D_6^{(1)}$, II*, RIMS Kokyuroku Bessatsu. **B5** (2008), 137–152.
- [17] T. Tsuda, K. Okamoto and H. Sakai, *Folding transformations of the Painlevé equations*, Math. Ann. **331** (2005), 713–738.